

Patient Information for Consent

Amir Salama

OS28 Revision Total Shoulder Replacement

Get more information and references at www.aboutmyhealth.org
Tell us how useful you found this document at www.patientfeedback.org



The Information Standard  Certified Member

eidohealthcare.com

EIDO
HEALTHCARE

UNITED KINGDOM

Information about COVID-19 (Coronavirus)

Hospitals have robust infection control procedures in place. However, you could still catch coronavirus either before you go to hospital or once you are there. If you have coronavirus at the time of your procedure, this could affect your recovery. It may increase your risk of pneumonia and in rare cases even death. The level of risk varies depending on factors such as age, weight, ethnicity and underlying health conditions. Your healthcare team may be able to tell you if these are higher or lower for you. Talk to your surgeon about the balance of risk between going ahead with your procedure and waiting until the pandemic is over (this could be many months).

Please visit <https://www.gov.uk/coronavirus> for up-to-date information.

Information about your procedure

Following the Covid-19 (coronavirus) pandemic, some operations have been delayed. As soon as the hospital confirms that it is safe, you will be offered a date. Your healthcare team can talk to you about the risks of having your procedure if you have coronavirus. It is then up to you to decide whether to go ahead or not. The benefits of the procedure, the alternatives and any complications that may happen are explained in this document. If you would rather delay or not have the procedure until you feel happy to go ahead, or if you want to cancel, tell your healthcare team.

If you decide to go ahead, you may need to self-isolate for a period of time beforehand (your healthcare team will confirm how long this will be). If you are not able to self-isolate, tell your healthcare team as soon as possible. You may need a coronavirus test a few days before the procedure. If your test is positive (meaning you have coronavirus), the procedure will be postponed until you have recovered.


Coronavirus spreads easily from person to person. The most common way that people catch it is by touching their face after they have touched a person or surface that has the virus on it. Wash your hands with alcoholic gel or soap and water when you enter the hospital, at regular intervals after that, and when you move from one part of the hospital to another.

Be aware of social distancing. Chairs and beds are spaced apart. If your healthcare team need to be close to you, they will wear personal protective equipment (PPE). If you can't hear what they are saying because of their PPE, ask them to repeat it until you can. You may not be allowed visitors, or your visiting may be restricted.

Your surgery is important and the hospital and health professionals looking after you are well equipped to perform it in a safe and clean environment. Guidance about coronavirus may change quickly your healthcare team will have the most up-to-date information.

What is a revision total shoulder replacement?

A revision total shoulder replacement is an operation to take out your old shoulder replacement and put in a new one.



A total shoulder replacement

Your surgeon has recommended a revision total shoulder replacement operation. However, it is your decision to go ahead with the operation or not.

This document will give you information about the benefits and risks to help you to make an informed decision. If you have any questions that this document does not answer, ask your surgeon or the healthcare team.

What causes a shoulder replacement to fail?

A shoulder replacement can fail for the following reasons.

- Wearing out of the artificial joint – This releases tiny bits of plastic that can cause the joint to come loose from the bone.
- Infection in your shoulder replacement – Infection can make your shoulder replacement come loose from the bone.
- Dislocation (coming out of joint) – You may decide to have an operation to try to prevent this from happening again.
- Fracture (break) around your shoulder replacement – This can happen if you fall heavily on your shoulder.

These problems can make it difficult for you to use your shoulder, and sometimes cause pain. Your surgeon will discuss with you why they have recommended a revision total shoulder replacement.

What are the benefits of surgery?

You should be able to move your shoulder more comfortably and do more of your normal activities.

Are there any alternatives to surgery?

Surgery is the only treatment for a shoulder replacement that is failing.

If your shoulder replacement is wearing out, your bones will become increasingly damaged over time. It is usually best to have the operation soon after your shoulder replacement starts to loosen, to reduce the risk of complications.

If you have an infection in your shoulder replacement, using antibiotics over the long term can sometimes prevent your shoulder replacement from failing. However, the infection will not settle unless your shoulder replacement is removed.

If your shoulder replacement keeps coming out of joint, you can wear a brace to try to keep your shoulder in place. A brace is large and often uncomfortable.

If you have a fracture around your shoulder replacement, it is likely that your surgeon will recommend an operation.

What will happen if I decide not to have the operation?

If your shoulder replacement is coming loose because of wear or infection, it will probably get more painful over time.

The bone around a loose shoulder replacement can get thin and it may break (fracture). You will need to have a major operation to fix the fracture and do your shoulder replacement again.

If you have an infection in your shoulder replacement, it can spread to other places around your body, including other artificial joints and vital organs.

Your surgeon will discuss with you what is likely to happen.

What does the operation involve?

The healthcare team will carry out a number of checks to make sure you have the operation you came in for and on the correct side. You can help by confirming to your surgeon and the healthcare team your name and the operation you are having.

Various anaesthetic techniques are possible. Your anaesthetist will discuss the options with you.

You may be given antibiotics during the operation to reduce the risk of infection.

Your surgeon will make a cut on the front of your shoulder.

If your shoulder is unstable and your shoulder replacement itself is not worn, your surgeon may leave your shoulder replacement and simply repair the ligaments around your shoulder or change some parts of your shoulder replacement.

If the joint surfaces have become worn or your shoulder replacement itself is coming loose, your surgeon will usually remove your shoulder replacement and any cement. This often takes a long time and can be difficult.

Your surgeon will put in a new shoulder replacement.

Your surgeon will close your skin with stitches or clips.

The type of surgery you need can be more complicated if the bone is thin or broken, or if you have an infection. Your surgeon will discuss with you what the operation is likely to involve.

- If your bone is thin or has broken, they may need to build it up with a bone graft, special wire mesh and cables, or a metal cage.

- If you have an infected shoulder replacement, you may need to have two separate operations. In the first, your surgeon will remove your old shoulder replacement, any cement and all infected material. They will put antibiotics in your shoulder and you will also need antibiotic injections for several weeks. An infection may take 2 to 3 months to go completely. You will need a second operation to put in the new shoulder replacement.

What should I do about my medication?

Make sure your healthcare team knows about all the medication you take and follow their advice. This includes all blood-thinning medication as well as herbal and complementary remedies, dietary supplements, and medication you can buy over the counter.

What can I do to help make the operation a success?

If you smoke, stopping smoking now may reduce your risk of developing complications and will improve your long-term health.

Try to maintain a healthy weight. You have a higher risk of developing complications if you are overweight.

Regular exercise should help to prepare you for the operation, help you to recover and improve your long-term health. Before you start exercising, ask the healthcare team or your GP for advice.

You can reduce your risk of infection in a surgical wound.

- In the week before the operation, do not shave or wax the area where a cut is likely to be made.
- Try to have a bath or shower either the day before or on the day of the operation.
- Keep warm around the time of the operation. Let the healthcare team know if you feel cold.

What complications can happen?

The healthcare team will try to reduce the risk of complications.

Any numbers which relate to risk are from studies of people who have had this operation. Your doctor may be able to tell you if the risk of a complication is higher or lower for you.

Some complications can be serious and can even cause death.

You should ask your doctor if there is anything you do not understand.

Your anaesthetist will be able to discuss with you the possible complications of having an anaesthetic.

General complications of any operation

- Pain. The healthcare team will give you medication to control the pain and it is important that you take it as you are told so you can move about as advised. Your surgeon may use a splint or sling to help to control the pain.
- Bleeding during or after the operation. You may need a blood transfusion.
- Infection of the surgical site (wound). It is usually safe to shower after 2 days but you should check with the healthcare team. Keep your wound dry and covered. Let the healthcare team know if you get a high temperature, notice pus in your wound, or if your wound becomes red, sore or painful. An infection usually settles with antibiotics but you may need another operation.
- Allergic reaction to the equipment, materials or medication. The healthcare team is trained to detect and treat any reactions that might happen. Let your doctor know if you have any allergies or if you have reacted to any medication or tests in the past.
- Unsightly scarring of your skin, although revision shoulder-replacement wounds usually heal to a neat scar.
- Blood clot in your leg (deep-vein thrombosis – DVT). This can cause pain, swelling or redness in your leg, or the veins near the surface of your leg to appear larger than normal. The healthcare team will assess your risk. They will encourage you to get out of bed soon after the operation and may give you injections, medication, or special stockings to wear. Let the healthcare team know straightaway if you think you might have a DVT.

- Blood clot in your lung (pulmonary embolus), if a blood clot moves through your bloodstream to your lungs. Let the healthcare team know straightaway if you become short of breath, feel pain in your chest or upper back, or if you cough up blood. If you are at home, call an ambulance or go immediately to your nearest Emergency department.

- Chest infection. You may need antibiotics and physiotherapy.

- Heart attack (where part of the heart muscle dies). A heart attack can sometimes cause death.

- Stroke (loss of brain function resulting from an interruption of the blood supply to your brain). A stroke can sometimes cause death.

Specific complications of this operation

- Damage to nerves around your shoulder, leading to weakness, numbness or pain in your shoulder or arm (risk: less than 1 in 100). This usually gets better but may be permanent.
- Infection, which can result in loosening and failure of your shoulder replacement (risk: less than 1 in 100). You will usually need one or more further operations to control the infection.
- Loosening without infection. You may need another operation to do your shoulder replacement again (risk: 1 in 20 over 8 years).
- Rotator-cuff tears. The rotator cuff is formed from four muscles and tendons that attach your arm to your shoulder blade. You may need surgery to repair any tears.
- Dislocation of your shoulder replacement (risk: less than 1 in 50 in the first 5 years). You may need another operation if it keeps on happening.
- Stiff shoulder. You should get more movement than you had before the operation but sometimes movement does not improve. It will never feel quite the same as a normal shoulder.

How soon will I recover?

In hospital

After the operation you will be transferred to the recovery area and then to the ward. You will usually have an x-ray to check the position of your shoulder replacement.

The physiotherapist will help you to start moving your shoulder, usually after 1 to 2 days. If your shoulder is unstable, or if your surgeon had to repair ligaments or your surgery was complicated, you may need to rest your shoulder for up to 6 weeks before you start physiotherapy.

The physiotherapist will teach you how to look after your new shoulder.

Keep your wound dry for 4 to 5 days, and use a waterproof dressing when you have a bath or shower.

The healthcare team will tell you if you need to have any stitches or clips removed, or dressings changed.

You should be able to go home after 2 to 3 days. However, your doctor may recommend that you stay a little longer.

If you are worried about anything, in hospital or at home, contact the healthcare team. They should be able to reassure you or identify and treat any complications.

Returning to normal activities

To reduce the risk of a blood clot, make sure you carefully follow the instructions of the healthcare team if you have been given medication or need to wear special stockings.

You will need to keep your arm in a sling to keep the tension away from your shoulder joint.

The healthcare team will tell you when you can return to normal activities. It often takes longer to recover from a revision shoulder replacement than your first shoulder replacement.

To reduce the risk of problems, it is important to look after your new shoulder as you are told.

Regular exercise should help you to return to normal activities as soon as possible. Before you start exercising, ask the healthcare team or your GP for advice.

Do not drive until you are confident about controlling your vehicle and always check your insurance policy and with your doctor.

The future

Do not play contact sports, use heavy tools such as a hammer, or do any activities that may involve you falling.

Most people make a good recovery and most revision total shoulder replacements work well. It is important to follow the advice the physiotherapist gives you about exercises to strengthen your shoulder muscles.

An artificial shoulder never feels quite the same as a normal shoulder and it is important to look after it in the long term.

A revision total shoulder replacement can fail with time if it wears out or the original problem comes back. You may need another operation (risk: 3 in 20 by 15 years after the operation).

Summary

If your original shoulder replacement fails, you can usually have another operation to do your shoulder replacement again. If this revision operation is successful, you should be able to continue many of your normal activities.

Surgery is usually safe and effective but complications can happen. You need to know about them to help you to make an informed decision about surgery. Knowing about them will also help to detect and treat any problems early.

Keep this information document. Use it to help you if you need to talk to the healthcare team.

Some information, such as risk and complication statistics, is taken from global studies and/or databases. Please ask your surgeon or doctor for more information about the risks that are specific to you.

This document is intended for information purposes only and should not replace advice that your relevant healthcare team would give you.

Acknowledgements

Reviewer: Lennard Funk (MSc, FRCS)

Illustrator: Medical Illustration Copyright © Medical-Artist.com